





# Comparing Effectiveness of Family Planning Methods

## More effective

Less than 1 pregnancy per 100 women in 1 year



## Less effective

About 30 pregnancies per 100 women in 1 year

## How to make your method more effective

**Implants, IUD, female sterilization:** After procedure, little or nothing to do or remember

**Vasectomy:** Use another method for first 3 months

**Injectables:** Get repeat injections on time

**Lactational amenorrhea method, LAM (for 6 months):** Breastfeed often, day and night

**Pills:** Take a pill each day

**Patch, ring:** Keep in place, change on time

**Condoms, diaphragm:** Use correctly every time you have sex

**Fertility awareness methods:** Abstain or use condoms on fertile days. Newest methods (Standard Days Method and TwoDay Method) may be easier to use.

**Withdrawal, spermicides:** Use correctly every time you have sex



### Sources:

Steiner MJ, Trussell J, Mehta N, Condon S, Subramaniam S, Bourne D. Communicating contraceptive effectiveness: a randomized controlled trial to inform a World Health Organization family planning handbook. *Am J Obstet Gynecol* 2006;195(1):85-91.

World Health Organization/Department of Reproductive Health and Research (WHO/RHR), Johns Hopkins Bloomberg School of Public Health (JHSPH)/Center for Communication Programs (CCP). *Family Planning: A Global Handbook for Providers*. Baltimore, MD and Geneva: CCP and WHO, 2007.

Trussell J. Choosing a contraceptive: efficacy, safety, and personal considerations. In: Hatcher RA, Trussell J, Stewart F, Nelson AL, Cates W Jr., Guest F, Kowal D, eds. *Contraceptive Technology, Nineteenth Revised Edition*. New York: Ardent Media, Inc., in press.